

ON THE NONNEGATIVITY OF THE COEFFICIENTS OF SOME POLYNOMIALS OCCURRING IN THE THEORY OF CUBICAL SPHERES

JAMES HAGLUND

RÉSUMÉ. Dans un de ces derniers articles portant sur les sphères cubiques, G. Heteyi a donné une preuve par récurrence du fait que les coefficients d’une certaine suite de polynômes sont non négatifs. À l’occasion du colloque SFCA’96, G. Heteyi a proposé de trouver une interprétation combinatoire de ces coefficients. Dans cette courte note, on démontre que certains résultats connus sur la théorie des tours impliquent que les dits coefficients comptent bien des permutations sujettes à certaines contraintes.

ABSTRACT. As part of his recent research into Cubical Spheres, G. Heteyi gave an inductive proof that a certain sequence of polynomials have nonnegative coefficients. In a poster session at the FPSAC’96 conference he asked for a combinatorial interpretation of these coefficients. In this short note we show that known results in rook theory imply that the coefficients count permutations subject to certain constraints.

At the recent FPSAC ’96 conference held at the University of Minnesota, G. Heteyi presented a poster session entitled “Invariants of Cubical Spheres” [Het1], a full length version of which appeared in this journal [Het2]. The following lemma was part of his session.

Lemma. *Let $\Phi_{a,b,c}(x)$ denote the formal power series $\sum_{k \geq 0} k^a(k+1)^b(k+2)^c x^k$, where a, b and c are natural numbers. Then the following hold.*

(i) *We have*

$$\Phi_{a,b,c}(x) = \frac{A_{a,b,c}(x)}{(1-x)^{a+b+c+1}}, \quad (1)$$

where $A_{a,b,c}(x)$ is a polynomial with integer coefficients, of degree at most $a+b+c$.

(ii) *If $b > 0$ then the degree of $A_{a,b,c}(x)$ is at most $a+b+c-1$.*

(iii) *If $b > 0$ or $c = 0$ then $A_{a,b,c}(x)$ has only nonnegative coefficients.*

Heteyi’s proof used induction and differential equations satisfied by $\Phi_{a,b,c}(x)$. As a challenge he asked for a combinatorial interpretation of part (iii) of his lemma. In this note we use rook theory to show that if $b > 0$ or $c = 0$, the coefficients of $A_{a,b,c}(t)$ count permutations subject to certain constraints.

Reçu le 15 août 1996 et, sous forme définitive, le 26 juin 1997.

If B is an n -column Ferrers board with column heights $h_1 \leq h_2 \leq \dots \leq h_n$, let $r_k(B)$ be the number of ways of placing k non-attacking rooks on B , and if $h_n \leq n$, let $t_k(B)$ be the number of ways of placing n non-attacking rooks on the $n \times n$ square chessboard containing B , with exactly k rooks on B . For more detailed descriptions of these objects see [Sta] and [Rio]. Our proof depends on the following theorem.

$$\sum_{k=0}^{\infty} x^k \prod_{i=1}^n (k + h_i - i + 1) = \frac{1}{(1-x)^{n+1}} \sum_{k=0}^n x^k t_{n-k}(B). \quad (2)$$

Eq. (2) is the $q = 1$ case of a result of Garsia and Remmel [GaRe, p. 259].

Let B be the board which starts off with c columns with heights $2, 3, \dots, c+1$, followed by b columns with heights $c+1, c+2, \dots, c+b$, followed by a columns with heights $c+b, c+b+1, \dots, c+b+a-1$. Then $h_{a+b+c} = a+b+c-1$, the left-hand side of (2) equals the left-hand side of (1), and part (iii) of the lemma follows. The condition $b > 0$ or $c = 0$ is necessary to ensure B has non-decreasing column heights.

To deduce part (ii) of the lemma, note that if $c > 0$, the first column of B has height 2, which implies $t_0(B) = t_1(B) = 0$, since any placement of $a+b+c$ rooks must have a rook in row 1 and a rook in row 2. Thus the degree of A is at most $a+b+c-2$. If $c = 0$ and $b > 0$, the first column of B has height 1 so we get $t_0(B) = 0$, with the degree of A being at most $a+b+c-1$.

Remarks. If $b = 0$, let B be the board which starts with a columns with heights $0, 1, \dots, a-1$, followed by c columns with heights $a+2, a+3, \dots, a+c+1$. In this case $h_{a+b+c} = a+b+c+1$, so the combinatorial definition of the t_k doesn't apply, but (2) still holds if we define the t_k via Riordan and Kaplansky's inclusion-exclusion identity [KaRi]

$$\sum_{k=0}^n k! r_{n-k}(B) (x-1)^{n-k} = \sum_{k=0}^n x^k t_k(B).$$

In [HOW] it is shown that a result of F. Brenti [Bre, Theorem 4.4.1, p. 43] implies that for any Ferrers board satisfying $h_n \leq n$, the hit polynomial $\sum_{k=0}^n t_k(B) x^k$ has only real zeros.

Résumé substantiel en français. Au colloque SFCA'96, qui a eu lieu à l'Université du Minnesota, G. Heteyi a présenté un article intitulé « Invariants des sphères cubiques » [Het1], dont une version complète apparaît dans cette revue. L'auteur y définit les polynômes $A_{a,b,c}(x)$ comme suit :

$$A_{a,b,c}(x) := (1-x)^{a+b+c+1} \sum_{k \geq 0} k^a (k+1)^b (k+2)^c x^k,$$

où a, b et c sont des nombres naturels. De plus, dans les cas $b > 0$ ou $c = 0$, G. Heteyi démontre par récurrence que les coefficients sont non négatifs. Par la suite, il propose de trouver une interprétation combinatoire à ces coefficients. Dans cet article, on utilise des résultats connus sur la théorie des tours pour montrer que les dits coefficients comptent bien des permutations sujettes à certaines contraintes.

Si B est un tableau de Ferrers ayant n colonnes d'hauteurs $h_1 \leq h_2 \leq \dots \leq h_n$, où $h_n \leq n$, soit $t_k(B)$ le nombre de façons de placer n tours non attaquantes sur un échiquier de taille $n \times n$ contenant B , avec exactement k tours sur B . Notre preuve dépend du théorème suivant, qui est le cas $q = 1$ d'un résultat de Garsia et Remmel :

$$(1-x)^{n+1} \sum_{k=0}^{\infty} x^k \prod_{i=1}^n (k + h_i - i + 1) = \sum_{k=0}^n x^k t_{n-k}(B).$$

Pour un choix approprié des valeurs h_i , le côté gauche de l'équation ci-dessus se réduit à $A_{a,b,c}(x)$ et le résultat en découle aisément.

REFERENCES

- [Bre] F. Brenti, *Unimodal, log-concave and Pólya frequency series in combinatorics*, Mem. Amer. Math. Soc. **89**, no. 413 (1989).
- [GaRe] A. M. Garsia and J. B. Remmel, *Q-counting rook configurations and a formula for Frobenius*, J. Combin. Theory Ser. A **41** (1986), 246–275.
- [HOW] J. Haglund, K. Ono and D. G. Wagner, *Theorems and conjectures involving rook polynomials with real roots*, Proceedings of the “Topics in Number Theory” Conference (University Park, PA, 1997) (G. E. Andrews and K. Ono, eds.), Kluwer Academic Publishing, To appear.
- [Het1] G. Hetyei, *Invariants of Cubical Spheres*, Proceedings of the Eighth International Conference on Formal Power Series and Algebraic Combinatorics, Minneapolis, Minnesota, USA, 1996, pp. 247–258.
- [Het2] G. Hetyei, *Invariants de complexes cubiques*, Ann. Sci. Math. Québec **20** (1996), 35–52.
- [KaRi] I. Kaplansky and J. Riordan, *The problem of the rooks and its applications*, Duke Math. J. **13** (1946), 259–268.
- [Rio] J. Riordan, *An Introduction to Combinatorial Analysis*, Wiley, New York, 1958.
- [Sta] R. Stanley, *Enumerative combinatorics, Vol. 1*, Wadsworth and Brooks, Monterey, 1986.

J. HAGLUND

DEPARTMENT OF MATHEMATICS

THE MASSACHUSETTS INSTITUTE OF TECHNOLOGY

CAMBRIDGE MA 02139

E-MAIL: haglund@math.mit.edu